

Short Term Analysis of the Functional and Radiological Outcome of Distal Femoral Fractures Fixed with Locking Compression Plate by Minimally Invasive Plate Osteosynthesis (MIPO) Technique

Sidheswar Baskey¹

Abstract

Background: Fractures of distal femur are very complex injuries and there is significant difficulty to manage. These injuries are severe and have a potential to produce long term disability. These fractures often are unstable and comminuted and tend to occur in elderly or multiply-injured patients.

The fractures of distal femur account for 7% of all femoral fractures. If Hip fractures are excluded, 31% of fractures involve the distal femur. The fractures involving distal 15 cm of femur including distal femoral metaphysis (supracondylar) and articular surface (intercondylar) are classified as distal femur fractures [1].

Distal femur fractures exhibits bimodal age distribution. In young adults it occurs due to high velocity trauma like road traffic accidents. These patients often sustain multiple and compound injuries. Older patients sustain distal femur fractures mostly due to trivial fall occurring in elderly osteoporotic bone.

In 1960's most of these fractures were treated conservatively and documented better outcome than operative treatment. But with the advent of newer implants and modern techniques, these fractures are best treated with surgical stabilization. The newer modalities of treatment include minimally invasive plate osteosynthesis (MIPO) and Less invasive skeletal stabilization (LISS).

Aim of this study: To evaluate twenty cases of distal femur fractures fixed with locking compression plate by minimally invasive plate osteosynthesis (MIPO) technique in the Department of Orthopaedic Surgery at SCB Medical College and Hospital, Odisha between July 2018 to November 2020.

To prospectively analyse the clinical and radiological outcome of the above procedure.

Keywords: Minimally Invasive Plate Osteosynthesis (MIPO), Distal Femoral Fractures, Radiological Outcome, Locking Compression Plate

Background

Fractures of distal femur are very complex injuries and there is significant difficulty to manage. These injuries are severe and have a potential to produce long term disability. These fractures often are unstable and comminuted and tend to occur in elderly or multiply-injured patients.

The fractures of distal femur account for 7% of all femoral fractures. If Hip fractures are excluded, 31% of fractures involve the distal femur. The fractures involving distal 15 cm of femur including distal femoral metaphysis (supracondylar) and articular surface (intercondylar) are classified as distal femur fractures [1].

Distal femur fractures exhibits bimodal age distribution. In young adults it occurs due to high velocity trauma like road traffic accidents. These patients often sustain multiple and compound injuries. Older patients sustain distal femur fractures mostly due to trivial fall

occurring in elderly osteoporotic bone.

In 1960's most of these fractures were treated conservatively and documented better outcome than operative treatment. But with the advent of newer implants and modern techniques, these fractures are best treated with surgical stabilization. The newer modalities of treatment include minimally invasive plate osteosynthesis (MIPO) and Less invasive skeletal stabilization (LISS).

Aim of this study: To evaluate twenty cases of distal femur fractures fixed with locking compression plate by minimally invasive plate osteosynthesis (MIPO) technique in the Department of Orthopaedic Surgery at SCB Medical College and Hospital, Odisha between July 2018 to November 2020.

To prospectively analyse the clinical and radiological outcome of the above procedure.

Materials & Methods

Study topic: A short term analysis of the functional and radiological outcome Of distal femoral fractures fixed with locking compression plate by minimally invasive plate osteosynthesis (MIPO) technique.

Study Design: Prospective study

Study Venue: Department of Orthopaedics, SCB Medical College & Hospital, Cuttack, Odisha 753007.

Study center: SCB Medical College & Hospital, Cuttack between



¹Department of Orthopaedic Surgery, SCB Medical College and Hospital, Cuttack, Odisha, India.

Address of Correspondence

Dr. Sidheswar Baskey
Department of Orthopaedic Surgery, SCB Medical College and Hospital, Cuttack, Odisha, India.
E-mail: stylishdoc@gmail.com

Received 19/05/2021; Reviewed 16/06/2021; Accepted 25/11/2021; Published 10/01/2022

© 2022 Odisha Journal of Orthopaedics & Trauma | Available on www.ojonline.com | DOI: 10.13107/ojot.2022.v43i01.032

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial-Share Alike 4.0 License (<http://creativecommons.org/licenses/by-nc-sa/4.0>) which allows others to remix, tweak, and build upon the work non-commercially as long as appropriate credit is given and the new creation are licensed under the identical terms.

July 2018 - November 2020.

Inclusion criteria

- Patients in the age group of above 18 years
- Distal femoral fractures – Müllers type A, C1 and C2
- < 2 weeks of injury
- Fractures reducible by indirect methods
- Grade I and II compound injuries (Gustillo Anderson)

Exclusion Criteria

- Fractures with grade III compound injuries
- Active infection
- Muller type B & C3 fractures
- Skeletal immaturity
- > 2 weeks of injury (fracture may not reducible by indirect methods)
- Periprosthetic fractures
- Comatosed patients
- Patients with risk of infections like on immune suppressants drugs
- Pathological fractures

Results

Criteria for clinical fracture union:

1. No pain/tenderness on weight bearing
2. No pain/tenderness on palpation/examination
3. Ability to walk/perform activities of daily living with no pain.

Criteria for radiological union: [32]

Grade	Callus Formation	Fracture line	Stage of union
1	Homogeneous bone structure	Obliterated	Achieved
2	Massive .Bone trabeculae crossing fracture line	Barely discernible	Achieved
3	Apparent. Bridging of fracture line	Discernible	Uncertain
4	Trace. No bridging of fracture line	Distinct	Not achieved
5	No callus formation	Distinct	Not achieved

Knee society score: [34]

The outcome analysis done using American knee society scoring system. The total score is 200 comprises of

- Knee score 100
- Function score 100

Outcome	Knee score (100)	Function score (100)	Total score (200)
Excellent	80-100	80-100	160-200
Good	70-79	70-79	140-159
Fair	60-69	60-69	120-139
Poor	< 60	< 60	< 120

Table VI: Outcomes

Observation and Results

The results were analysed prospectively both clinically and radiologically. The follow up period ranged from 6 months to 18 months (mean = 12 month). The results of our study are elucidated

s. no	Age (in years)	No. of patients	Percentage
1	11-20	1	5
2	21-30	4	20
3	31-40	4	20
4	41-50	1	5
5	51-60	3	15
6	61-70	6	30
7	71-80	1	5

S. no	Sex	No. of patients	Percentage
1	Male	13	65
2	Female	7	35

S. no	Hammer et al Grade	No. of patients	Percentage
1	I	12	60
2	II	8	40

Graph 1: Hammer et al Grading






Table IV: Knee Extension

S. no	KNEE EXTENSION	No. of patients	Percentage
1	Full	17	85
2	Extensor lag	3	15



Graph 2: Knee Extension



Follow up	Minimum flexion	Maximum flexion	Mean flexion
1 month	10	90	56.5
3 month	20	105	70.25
6 month	20	120	84.25
Final	30	125	91.75



Follow up	Minimum score	Maximum score	Mean score
1 month	32	78	57
3 month	56	146	111
6 month	68	166	133
Final	81	188	153



Author	No. of cases	F/up months mean	Mean Time to union(wks)	Knee Flexion deg	complications			scoring system & mean score	excellent and good results %
					Infection %	Revision/Failure %	Malalign %		
Kregor et al	66	9	11	103	3	1.5	4.5	-	-
Kanabar et al	17	12	17	93	-	11.7	6	-	-
Schutz et al	99	13.7	-	107	7	6	1	-	-
Markmiller et al	20	12	13.8	110	-	10	15	Lysholm	87.5
Schandelmaier et al	54	6	14.3	104	1.4	7.9	13	Neer	-
Yeap & Deepak et al	11	9.7	18	107	-	9	9	schatzker	72.7
our study	20	12	15	92	5	5	-	Kss Mean-153	75

in the tables presented below. Table I, II, III, IV, V, VI, VII, VIII, IX and graphs 1, 2 & 3.

Discussion

The evolution of management of distal femoral fractures has come a long way from totally conservative management in the 1960's to definitive surgical treatment at present. There is increasing incidence of comminuted distal femur fractures due high velocity motor vehicle accidents in younger population and increased life expectancy resulting in fractures following trivial fall due to osteoporotic bone. In both the groups our aim is to restore the function and near normal anatomy similar to the pre injury status.

There are many surgical alternatives for distal femur fractures, each with its own pearls and pitfalls. Many studies were conducted using different implants and techniques resulted in varying outcome and complications. After the introduction of locking compression plate (LCP) by AO in 2000, the trend is shifting towards it due its added advantages like,

- Providing both angular and axial stability
- Applied in both locking and compression mode
- Better hold in osteoporotic bone

Due to the changing concepts towards relative stability and biological fixation from absolute stability and rigid fixation, minimally invasive plate osteosynthesis (MIPO) technique evolved. Many studies proved better outcome with lesser morbidity than the conventional technique.

In our study involving 20 patients with 13 males and 7 females with mean age of 47 and the mean follow up period ranges between 6 month to 18 month (mean 12 months). 15 patients had type A fracture and 5 patients had type C fracture and 25% of patients had open injuries.

The mean operating time was 100 minutes compared to 119 minutes by Yeap and Deepak et al [35]. The mean time to radiological fracture union was 15 weeks (range 12 -20 weeks) Which was comparable to 11 weeks by Kregor et al [30], 14.3 weeks by Schandelmaier et al [28], 12 weeks by Fankhauser et al [34] and 18 weeks by Yeap and Deepak et al [35].

The average knee flexion achieved was 92 degrees comparable to that of 103° by Kregor et al [30], 104° by Schandelmaier et al [28], 107° by schutz et al [29], 101° by Fankhauser et al [34] and 93° by Kanabar et al [36].

The scoring system used was knee society scoring and the mean score was 153 compared to the score of 131 by Fankhauser et al . With this system 45% patient is having an excellent outcome, 30% good, 15% fair and 10% with poor outcome.

The percentage of patients with good and excellent outcome was 75% comparable to 87.5% by Markmiller et al, 72.7% by Yeap and Deepak et al.

The complications encountered are deep seated infection (n=1), post-operative loss of reduction which requires a revision surgery (n=1), deep vein thrombosis (n=1), Knee stiffness (n=3), varus malalignment (n=2), reactive synovitis (n=1).

The incidence of loss of reduction requiring a revision surgery was 5% comparable to 10 % by Markmiller et al, 9 % by Yeap and Deepak et al 7.9 % by Schandelmaier et al and 6 % by Schutz et al.

The infection rate in our study was 5 % comparable to 7 % by Schutz

et al [29] and 3% by Kregor et al [30].

We had two cases of varus malalignment ($< 5^\circ$) but within acceptable limits in contrast to 15% by Markmiller et al and 13% by Schandelmaier et al, both having significant malalignment ($> 10^\circ$). These patients may require a long term follow up to evaluate the development of arthritis.

There was an incidence of 15% knee stiffness ($n=2$) $< 30^\circ$ and ($n=1$) 70° and failed to show any improvement even after aggressive continuous motion therapy.

We had a complication of post op DVT and reactive synovitis, which settled uneventfully with symptomatic therapy. Union was achieved in all cases and bone grafting was not required in any of our cases.

Conclusion

- In our study, Minimally Invasive Plate Osteosynthesis (MIPO) technique using Locking Compression Plate (LCP) shows good to excellent results in terms of union and functional outcome.
- When operated within two weeks of injury, it was easier to achieve

closed reduction. This decreases the operating time, blood loss and intra-operative morbidity.

- MIPO technique could result in satisfactory union and eliminates the need for bone grafting.
- The incidence of infection and post-operative morbidity was less compared to conventional open technique.
- LCP has a better hold in osteoporotic bone with less chances of failure.
- Inadequate fixation leads to loss of reduction, resulting in an open reduction and revision fixation.
- Long term follow up is necessary to study the development of arthritis in patients with varus/valgus malalignment.
- From our study, we conclude that the Minimally Invasive Plate Osteosynthesis (MIPO) technique using Locking Compression Plate (LCP) will result in early post-operative rehabilitation, satisfactory union and good functional outcome. The chances of infection and implant failure are less. Proper patient selection and meticulous surgical techniques will give the best results.

References

1. Charles A. Rockwood RWB. Rockwood and Green's Fractures in Adults. 17th ed.: Lippincott Williams & Wilkins; 2010.
2. Snell RS. clinical anatomy by regions. 9th ed.: Lippincott Williams & Wilkins; 2011.
3. Henry Gray S. Gray's Anatomy: The Anatomical Basis of Clinical Practice: Elsevier Science Health Science Division; 2005.
4. By A. M. R. Agur AFDJCBG. Grant's atlas of Anatomy. 13th ed.: Lippincott Williams & Wilkins; 2012.
5. AS Reddy RF. Evaluation of the Intraosseous and Extraosseous Blood Supply to the Distal Femoral Condyles. American journal of sports medicine. may 1998.
6. Reto Babst SB. AO manual Minimally Invasive Plate Osteosynthesis (MIPO); 2012.
7. S. Pickering DA. Alignment in Total Knee Replacement. bone & joint journal..
8. Donald R. Peterson JDB. Biomechanics: Principles and Applications, Second Edition.
9. S.Terry canale JHB. Campbell's operative orthopaedics. 12th ed.: elsevier; 2013.
10. Stewart MJ ST,WS. Fractures of the distal third of femur. J Bone and joint surgery Am. 1966;(48): 784-807).
11. NEERII CS, GRANTHAM SA, SHELTON ML. Supracondylar Fracture of the Adult Femur a study of one hundred and ten cases. J Bone Joint Surg Am, 1967 Jun 01;49(4):591-613. 1967 june.
12. J Schatzker GHJW. The Toronto experience with the supracondylar fracture of the femur, 1966–1972. Injury. 1974, 6(2):113-128.
13. al BBe. The results of open reduction and Internal fixation of distal femur fractures using a biologic (indirect) reduction technique. J orthop Trauma. 1996;10(6):372-7..
14. Ritter MA,PMFaEMK. "Anterior femoral notching and ipsilateral supracondylar femur fracture in total knee arthroplasty." The Journal of arthroplasty 3.2 (1988): 185-187..
15. Yuvarajan RKaLM. "Review of concepts in distal femoral fractures management." Pb Journal of orthopaedics 11.1 (2009)..
16. al BBe. The use of modified Neufeld traction in the management of femoral fractures in polytrauma. journal of trauma: 1981 Sep;21(9):779-87..
17. MOONEY Vea. "Cast-Brace Treatment for Fractures of the Distal Part of the Femur a prospective controlled study of one hundred and fifty patients. The Journal of Bone & Joint Surgery 52.8 (1970): 1563-1578..
18. Butt MS,SJKaMSA. Displaced fractures of the distal femur in elderly patients. JOURNAL OF BONE AND JOINT SURGERY-BRITISH VOLUME-78 (1996): 110-118..
19. Michael Wagner RF. AO manual of Internal fixator - concepts and cases using LISS and LCP: Thieme; 2012.
20. Henry SL,ea. "Management of supracondylar fractures of the femur with the GSH intramedullary nail: preliminary report." Contemporary orthopaedics 22.6 (1991): 631..
21. Ricci WMea. Angular Malalignment After Intramedullary Nailing of Femoral Shaft Fractures. Journal of Orthopaedic Trauma: 2001; 15(2) 90-95..
22. Thomson ABea. "Long-term functional outcomes after intra-articular distal femur fractures: ORIF versus retrograde intramedullary nailing." Orthopedics 31.8 (2008): 748-750..
23. Fixation ToDFaPTFWEFFbPCTl. Parekh, Anand A.; Smith, Wade R.; Silva. Journal of Trauma-Injury Infection & Critical Care. 64(3):736-739, March 2008..
24. Arazi M,ea. "Ilizarov external fixation for severely comminuted supracondylar and intercondylar fractures of the distal femur." Journal of Bone & Joint Surgery, British Volume 83.5 (2001): 663-667..
25. al Ee. Biomechanics of locked plates and screws. J orthop Trauma. ; 18(8):483-487.
26. al WMe. Treatment of distal femoral fractures in the elderly using less invasive plating technique. Int orthop. ; 29(2):117-120.

27. Mast JJRGR. Planning and Reduction Technique in Fracture Surgery. 1st ed.: springer; 1989.
28. Schandelmaier P,ea. "Distal femoral fractures and LISS stabilization." *Injury* 32 (2001): 55-63..
29. Schütz M,ea. "Less invasive stabilization system (LISS) in the treatment of distal femoral fractures.". *Acta chirurgiae orthopaedicae et traumatologiae Cechoslovaca* 70.2 (2003): 74..
30. Kregor PJ,ea. "Distal femoral fracture fixation utilizing the Less Invasive Stabilization System (LISS): the technique and early results." *Injury* 32 (2001): 32-47..
31. Kim SJ,ea. "Minimally invasive plate osteosynthesis for distal femoral fractures." *Journal of the Korean Society of Fractures* 16.4 (2003): 474-481..
32. Whelan DB,ea. "Interobserver and intraobserver variation in the assessment of the healing of tibial fractures after intramedullary fixation." *Journal of Bone & Joint Surgery, British Volume* 84.1 (2002): 15-18..
33. KILIÇOĞLU ÖI,ea. "Comparison of locked plating and intramedullary nailing for periprosthetic supracondylar femur fractures after knee arthroplasty." *Acta orthopaedica Belgica* 79.4 (2013): 417-421..
34. Fankhauser Fea. "Minimal-invasive treatment of distal femoral fractures with the LISS (Less Invasive Stabilization System) A prospective study of 30 fractures with a follow up of 20 months." *Acta Orthopaedica* 75.1 (2004): 56-60..
35. Yeap EJ,aASD. "Distal femoral Locking Compression Plate fixation in distal femoral fractures: early results." *Malaysian Orthop J* 1.1 (2007): 12-17..
36. Kanabar P,ea. "Less invasive stabilisation system plating for distal femoral fractures." *Journal of Orthopaedic Surgery* 15.3 (2007).

Declaration of patient consent: The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient has given his/her consent for his/her images and other clinical information to be reported in the Journal. The patient understands that his/her name and initials will not be published, and due efforts will be made to conceal his/her identity, but anonymity cannot be guaranteed.

Conflict of Interest: NIL
Source of Support: NIL

How to Cite this Article

Baskey S | Short Term Analysis of the Functional and Radiological Outcome of Distal Femoral Fractures Fixed with Locking Compression Plate by Minimally Invasive Plate Osteosynthesis (MIPO) Technique | *Odisha Journal of Orthopaedics & Trauma* | January 2022; 43: 39-43.